



University of Groningen

STAT5-mediated self-renewal of normal hematopoietic and leukemic stem cells

Schepers, Hein; Wierenga, Albertus T. J.; Vellenga, Edo; Schuringa, Jan Jacob

Published in:
JAK-STAT

DOI:
[10.4161/jkst.19316](https://doi.org/10.4161/jkst.19316)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2012

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Schepers, H., Wierenga, A. T. J., Vellenga, E., & Schuringa, J. J. (2012). STAT5-mediated self-renewal of normal hematopoietic and leukemic stem cells. *JAK-STAT*, 1(1), 13-22. <https://doi.org/10.4161/jkst.19316>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

STAT5-mediated self-renewal of normal hematopoietic and leukemic stem cells

Hein Schepers,^{1,2} Albertus T.J. Wierenga,^{1,3} Edo Vellenga¹ and Jan Jacob Schuringa^{1,*}

¹Department of Experimental Hematology; University Medical Center Groningen; Groningen, The Netherlands; ²Department of Stem Cell Biology; University Medical Center Groningen; Groningen, The Netherlands; ³Department of Laboratory Medicine; University Medical Center Groningen; Groningen, The Netherlands

The level of transcription factor activity critically regulates cell fate decisions such as hematopoietic stem cell self-renewal and differentiation. The balance between hematopoietic stem cell self-renewal and differentiation needs to be tightly controlled, as a shift toward differentiation might exhaust the stem cell pool, while a shift toward self-renewal might mark the onset of leukemic transformation. A number of transcription factors have been proposed to be critically involved in governing stem cell fate and lineage commitment, such as Hox transcription factors, c-Myc, Notch1, β -Catenin, C/ebp α , Pu.1 and STAT5. It is therefore no surprise that dysregulation of these transcription factors can also contribute to the development of leukemias. This review will discuss the role of STAT5 in both normal and leukemic hematopoietic stem cells as well as mechanisms by which STAT5 might contribute to the development of human leukemias.

Introduction

Signal Transducer and Activator of Transcription 5 (STAT5) is widely expressed throughout the hematopoietic system, both in stem and progenitor cells as well as in committed erythroid, myeloid and lymphoid cells.^{1–3} Indeed, it is not surprising that STAT5 can be activated by a wide variety of cytokines and growth factors.^{3–5} These include cytokines and growth factors that can signal through the Interleukin 3 (IL3)-receptor family [IL3, IL5, Granulocyte-Macrophage Colony Stimulating Factor (GM-CSF)], through the common γ -chain receptor family (IL2, IL7, IL9, IL12, IL15), through single chain receptors [Erythropoietin (EPO), Thrombopoietin (TPO), Growth Hormone (GH), prolactin, Granulocyte-Colony Stimulating Factor (G-CSF)], through class II receptors [Interferon α (IFN α), IFN γ , IL22] or through tyrosine kinase receptors [Stem Cell Factor (SCF), Platelet Derived Growth Factor (PDGF), Epidermal growth Factor (EGF)] (Fig. 1). In most cases, Janus Kinase (JAK) tyrosine kinase activity mediates STAT5 tyrosine phosphorylation, and STAT5 can be activated by JAK1, 2, or 3, depending on the cytokine-activated receptor complex. Alternatively, the tyrosine kinase receptor family can also induce STAT5 phosphorylation in a JAK-independent manner. While STAT5 is expressed in the majority of hematopoietic cell types, the cytokine receptor expression is much more tissue-specific. Thus, specific cytokines

are able to induce STAT5 activity in subsets of cell types only.⁶ For example, cytokines that activate STAT5 in the most immature human hematopoietic stem compartment include SCF⁷ and TPO.⁸ These cytokines have been shown to promote long-term hematopoiesis *in vitro*,⁹ and hypersensitivity to TPO in Lnk^{-/-} mice resulted in elevated stem cell self-renewal, which coincided with increased levels of STAT5 activity.¹⁰ Within the erythroid compartment, STAT5 is activated by EPO,¹¹ where STAT5 fulfills an important anti-apoptotic role by upregulating Bcl-X_L,^{12–15} although a more direct role in initiating erythroid commitment might exist as well.^{16–18} In myeloid cells, STAT5 can be activated by a variety of cytokines, including IL3, IL5, GM-CSF and CSF1 (reviewed in ref. 3). Although initially in STAT5AB^{AN/AN} mice myelopoiesis appeared to be relatively unaffected,¹⁹ it is likely that in myeloid cells many of the signals initiated by e.g., IL-3 and GM-CSF are, at least in part, mediated by STAT5,^{20,21} thereby regulating myeloproliferation or anti-apoptosis.^{22–24} During myelosuppression, mice completely deficient of STAT5AB failed to produce enhanced levels of neutrophils and were unable to respond to GM-CSF.²⁵ IL5-induced STAT5 activation is required for the induction of eosinophil differentiation.²⁶ Lymphoid development is severely impaired in STAT5^{-/-} mice.²⁷ STAT5 activation is required for IL2-induced T cell proliferation and the production of NK cells,^{28,29} or for IL7-mediated B cell expansion.³⁰

As summarized in Table 1 and Figure 1B, a wide variety of genetic defects in myeloid leukemias and myeloproliferative diseases (MPDs) result in activation of the STAT5 pathway, including mutations in Flt3 and cKit receptors, JAK2 mutations, translocations such as TEL-PDGFRA, and Bcr-Abl, but also as a result of increased cytokine signaling. Numerous functional studies have indicated that aberrant activation of STAT5 can contribute to the process of leukemic transformation. Downstream of Flt3-ITD (Internal Tandem Duplication) mutations, STAT5 is strongly activated via two tyrosine residues within the Flt3 receptor, Y589 and Y592 that act as docking sites for the SH2 domain of STAT5 molecules.³¹ Mutation of these residues into phenylalanines completely abrogated activation of STAT5, and importantly completely impaired induction of a myeloproliferative disease *in vivo* in a murine transplantation model.³¹ Thus, it is likely that STAT5 signaling is essential for the transforming potential of Flt3-ITD.

Although in human cells introduction of Flt3-ITD did not result in a myeloproliferative disease in transplanted NOD-SCID mice, the activated stem cell phenotype imposed on CB CD34⁺ cells, as revealed by the formation of early cobblestone area

*Correspondence to: Jan Jacob Schuringa; Email: j.schuringa@int.umcg.nl
Submitted: 11/29/11; Revised: 01/10/12; Accepted: 01/11/12
<http://dx.doi.org/10.4161/jkst.1.1.19316>




Figure 1. STAT5 signaling in normal and leukemic cells. (A) Normal cytokine-induced STAT5 signaling. (B) Constitutive STAT5 signaling in hematological malignancies.

forming cells (CAFCs), was impaired by coexpression of a dominant negative STAT5A (Y694F) mutant,³² suggesting that also in human cells STAT5 is an important mediator of Flt3-ITD-induced signaling. In studies in which STAT5 expression was targeted in primary acute myeloid leukemia (AML) CD34⁺ cells using a lentiviral approach, it was observed that long-term expansion and the formation of leukemic CAFCs was strongly impaired by downmodulation of STAT5.³³ Although the presence of Flt3-ITDs was not the exclusive genetic mutation that induced constitutive STAT5 signaling in the samples that were studied,

these data clearly underscore the important role that STAT5 fulfills in long-term expansion and self-renewal of primary AML stem/progenitor cells as well.

In chronic myeloid leukemia (CML) induced by Bcr-Abl, it has been convincingly demonstrated that STAT5 also plays an important role. A number of studies have shown that STAT5 is efficiently activated downstream of Bcr-Abl,⁵⁹⁻⁶² and interference with STAT5 activation negatively impacts the survival and proliferation of Bcr-Abl-expressing cells.^{27,55,63-69} Bcr-Abl p210-transduced STAT5^{AN/AN} murine bone marrow (BM) cells developed CML with low frequencies, with a delayed onset of disease.⁵⁵ The induction of acute lymphoid leukemia (ALL) was not impaired in these animals. Complete abrogation of STAT5 expression in STAT5AB^{-/-} mice also impaired lymphoid transformation induced by Bcr-Abl-expressing murine BM.^{27,70} In primary human CML cells, it was demonstrated that downmodulation of STAT5 expression by RNAi impaired Bcr-Abl-dependent proliferation and also reduced colony formation in methylcellulose.⁶⁸ Inhibition of STAT5 by pimozide reduced colony formation of CML CD34⁺ cells, also in tyrosine kinase-resistant patient samples.⁷¹

In myeloproliferative diseases it has been demonstrated in mouse models that bone marrow (BM)-transduced with TEL-JAK2 no longer induced disease in recipient mice when the oncogene was introduced in a STAT5^{AN/AN} background.⁴⁰ Similarly, MPD induced by TEL-PDGFR α depended on STAT5 activity.^{50,51} Finally, enhanced STAT5 activity has been observed in Polycythemia Vera (PV), caused by the activating JAK2 V617F mutation.⁷² Inhibition of JAK2 kinase activity abrogated the activation of STAT5, which coincided with a suppression of erythropoiesis in vitro and in vivo.^{73,74}

The most direct evidence for STAT5 acting as an oncogene arises from murine BM transplantation studies in which constitutively activated STAT5 (S711F) mutants were overexpressed. Lethally irradiated recipients receiving activated STAT5-transduced BM died within 6 weeks after transplantation of a multilineage leukemia.⁷⁵ It was demonstrated that a tryptophan residue in the N-terminal region of STAT5 is required for tetramerization of STAT5 dimers, and tetramer-deficient STAT5 mutants were unable to induce leukemia in mice.⁷⁵ Another activating mutant of STAT5, STAT5A(1*6) that contains two point mutations (H299R and S711F)⁷⁶ was earlier shown to induce myeloid hyperproliferation, but not leukemia, in a murine

Table 1. STAT5 activation in hematological malignancies

Kinase	Mutation/Translocation	Disease	References
Flt3	ITD (exon 11–12, 3–400 bp) TK point mutations (Asp835)	AML	31,34–38
		AML	39
JAK2	TEL-JAK2 V617F	ALL, CML, MPD	40
		PV, ET, IMF	41–44
JAK3	A572V, V722I, P132T	AMKL	45
c-Kit	D816V ITD (exon 11–12)	AML	46–48
		Pediatric AML	49
PDGFR	TEL-PDGFR FIP1L1-PDGFR α	CML, MPD	50,51
		Chronic Eosinophilic Leukemia	52,53
Abl	v-ABL (p160) BCR-ABL (p210, p185)	CML, BCL	54
		CML	27,55
FGFR	ZNF198-FGFR1	MPD	56
RAR α	STAT5b-RAR α	APL	57,58

retroviral overexpression model.⁴⁰ This was later confirmed by others, and a fatal MPD was observed by overexpression of these STAT5 mutants, but only when the most primitive CD34⁺Lin[−]cKit⁺Sca1⁺ (LSK) population was transduced and used for transplantation to irradiated recipients, suggesting that the stem cell, but not a committed progenitor is the target cell for transformation induced by activated STAT5.⁷⁷ Intriguingly, while these examples clearly demonstrate that STAT5 can transform murine hematopoietic stem cells (HSCs), no in vivo STAT5-induced transformation has been reported in human cell populations. While enhanced self-renewal and long-term stem cell maintenance can be achieved by introduction of activated STAT5 in human CD34⁺ cells, a myeloproliferative disease or leukemia does not occur in non-obese diabetic/severe combined immunodeficiency (NOD-SCID) transplantations models.¹⁶ It is plausible that the NOD-SCID xenograft model is not suitable to completely recapitulate human disease, or alternatively it is possible that species-specific differences in STAT5 signaling exist. In line with these observations, introduction of Bcr-Abl in murine BM resulted in a rapid and lethal MPD whereby recipients die within 3 weeks after transplantation. Introduction of Bcr-Abl in human CD34⁺ cells does not result in a rapid leukemia or MPD in engrafted NOD-SCID mice, and only after 5 mo progression to an early stage disease was observed in some animals.^{78,79}

Collectively, these data indicate that STAT5 is frequently activated in various hematological malignancies, whereby it strongly affects processes such as self-renewal and lineage fate determination. Whether STAT5 target genes in normal and leukemic stem cells are identical, or whether leukemic stem cell-specific STAT5 target genes exist remains to be determined. Also, it will be informative to study how STAT5 might cooperate with additional leukemic oncogenes in a multi-hit approach to model the development of human leukemias.

STAT5 as a Stem Cell Self-Renewal Factor

Loss-of-function and gain-of-function experiments have revealed critical roles for STAT5 in the hematopoietic stem/progenitor

compartment. STAT5AB^{AN/AN} mice have been used to assess stem cell function in the absence of wt STAT5 signaling. These mice were characterized by normal HSC numbers and stem cells isolated from the bone marrow or fetal liver were capable of engrafting irradiated recipients.⁸⁰ Yet, competitive repopulating capacity of STAT5AB^{AN/AN} HSCs was severely impaired.^{80–85} The underlying mechanisms are not fully elucidated yet, but it has been observed that the responsiveness of STAT5AB^{AN/AN} HSCs to early-acting cytokines such as IL3 and SCF was reduced, while the sensitivity to 5-fluorouracil was enhanced.⁸¹ Loss of protection against apoptosis most likely does not explain the STAT5AB^{AN/AN} HSCs phenotypes, as overexpression of Bcl2 was not sufficient to rescue repopulating defects.⁸⁴ Although homing of STAT5AB^{AN/AN} BM cells into lethally irradiated recipients was not impaired,⁸³ retention in the bone marrow was reduced under non-myeloablative conditions,⁸⁶ leaving open the possibility that competition for the niche might play a role. Mice completely devoid of STAT5 display severely impaired hematopoiesis.⁸⁷ More recently, using an Mx1-Cre inducible mouse model, it was shown that conditional deletion of STAT5 results in a loss of stem cell quiescence, associated with reduced survival and gradual loss of the HSC pool.⁸⁸

In order to study STAT5 signaling in human hematopoietic stem/progenitor cells, we have used a lentiviral shRNA approach in cord blood (CB) CD34⁺ cells.³³ Downmodulation of STAT5 to about 30% of the endogenous levels reduced progenitor frequencies as determined by Colony Forming Cell (CFC) assays in methylcellulose as well as stem cell frequencies as determined by Long-Term Culture-Initiating Cell (LTC-IC) assays in limiting dilution.³³ This resulted in reduced long-term expansion on MS5 bone marrow stroma upon downmodulation of STAT5 expression, whereby the myeloid and erythroid differentiation were unaffected. Single-cell assays using transduced CD34⁺/CD38[−] cells revealed that cell cycle progression induced by early-acting cytokines SCF and TPO was impaired by STAT5 downmodulation.³³

Reversely, activating mutants of STAT5 have been introduced in murine CD34⁺LSK cells and the effects on stem and progenitor cells were assessed in vitro and in vivo.⁷⁷ Introduction of STAT5A(1*6) or STAT5A(1*7) mutants resulted in a strong ex vivo expansion of immature CFU-nmEM progenitors, without affecting the symmetry of stem cell divisions as determined in paired-daughter cell assays.⁷⁷ Importantly, long-term repopulating HSCs could be maintained under ex vivo culture conditions as CD34⁺LSK cells expressing activated STAT5 had a strong competitive repopulating advantage over wild type cells after 7 d and 10 d ex vivo culturing in the presence of SCF or SCF and TPO.⁷⁷

Activating mutants of STAT5 have also been introduced in human CD34⁺ cells. Overexpression of STAT5A(1*6) in human CD34⁺ cord blood cells resulted in enhanced stem cell self-renewal.¹⁶ This enhanced self-renewal was only observed in bone marrow stromal cocultures, but not in cytokine-driven liquid culture conditions. These data argued that STAT5-induced HSC cell self-renewal depends on the presence of a bone marrow microenvironment, and it was indeed observed that STAT5A(1*6)-expressing CD34⁺ cells have a strongly enhanced interaction

with bone marrow stromal cells, resulting in the appearance of early CAFCs underneath the stroma within 1 week after plating. These CAFCs contained self-renewal potential as demonstrated by their capacity to give rise to second CAFCs upon harvest and replating onto new stroma, as well as by their capacity to engraft in sublethally irradiated NOD-SCID mice. Upon serial replating, long-term cultures could be established by overexpression of activated STAT5 for over 20 weeks, giving rise to new CAFCs upon each replating as well as to progeny in suspension. Hematopoietic progenitors could be maintained long-term in these culture conditions and the suspension cells retained an immature blast-like morphology.¹⁶

When STAT5A(1*6) mutants were expressed in murine embryonic stem (ES) cells, the generation of hematopoietic stem cells was greatly facilitated as studied on OP9 bone marrow stromal cells.¹⁷ The generation of hematopoietic CAFCs was strongly enhanced by activated STAT5. Importantly, these CAFCs could be serially passaged onto new OP9 stroma, giving rise to second and third CAFCs that were able to sustain long-term hematopoiesis and generate high numbers hematopoietic progenitors, indicative of HSC self-renewal in vitro. Also, the CAFCs generated by activation of STAT5 could engraft sublethally irradiated NOD-SCID mice, indicating that STAT5 facilitates the generation of ES-derived HSCs that can contribute to hematopoiesis in vivo as well.¹⁷

Mechanisms Involved in STAT5-Induced HSC Self-Renewal

Although various STAT5 target genes have been identified, the mechanisms by which STAT5 acts on HSCs remain to be elucidated. Using cell lines or heterogeneous stem/progenitor cell populations, enhanced cell growth is one of the most dominant phenotypes that is frequently observed in various studies, and several genes that are regulated by STAT5 associate with cell proliferation and cell cycle progression, including Cyclin D1, Pim1, and c-Myc.^{16,89-91} CyclinD1^{-/-}D2^{-/-} mice exhibit defects in the expansion of hematopoietic stem/progenitor cells.⁹² Pim serine/threonine kinases act as mediators of cytokine-induced cell growth⁹³ by promoting acceleration of cell-cycle progression both at the G1/S and G2/M transitions by phosphorylating and activating the phosphatases Cdc25A and Cdc25C, respectively.^{94,95} c-Myc controls the balance between self-renewal and differentiation of HSCs.^{96,97} However, when more purified stem cell populations were studied using a conditional deletion Mx1-Cre model it was observed that STAT5 was required to maintain HSC quiescence. Upon STAT5 deletion a decrease in the percentage of cells in G0 within the long-term and short-term HSC compartments was observed, coinciding with a decrease in expression of quiescence-associated genes such as p57 and Tie2.⁸⁸ Loss of STAT5 also coincided with an increase in AnnexinV⁺ CD34⁺ LSK cells.⁸⁸ In erythroid cells, it has been convincingly demonstrated that STAT5 contributes to cell survival by upregulating the anti-apoptosis gene Bcl-X_L.^{13,14} Survival of Flt3-ITD⁺ AML cells has been shown to depend on STAT5-mediated expression of Mcl1.⁹⁸ Also, Bcl2 can be upregulated by

STAT5 and is required to prevent apoptosis during terminal differentiation of myeloid cells.²³ Whether prevention of apoptosis contributes to STAT5-induced HSC self-renewal is currently unclear, but Bcl2 overexpression was not sufficient to rescue the repopulation defects of STAT5A^{AN/AN} HSCs,⁸⁴ suggesting that protection against apoptosis is not the main role of STAT5 signaling in HSCs. In our co-cultures, despite strong reductions in LTC-IC and CFC frequencies, we also did not detect an increased rate of apoptosis in STAT5 RNAi-transduced CD34⁺ CB cells, and no decreased expression of the Bcl-X_L gene was observed.³³ The basic helix-loop-helix transcriptional inhibitor ID1 is also upregulated by STAT5,⁹⁹ and ID1-deficient HSCs fail to self-renew, leading to low steady-state HSC numbers and premature HSC exhaustion.¹⁰⁰ Little evidence exists that STAT5 affects the expression of other known HSC self-renewal regulators such as Bmi1 or HoxB4. Recently, we observed that STAT5 binds to and activates the promoter of Hypoxia Induced Factor 2 α (Hif2 α) in human CD34⁺/CD38⁻ HSCs.¹⁰¹ Functional studies indicated that STAT5-induced long-term expansion and elevated LTC-IC and CFC frequencies were reduced upon downmodulation of Hif2 α .¹⁰¹ Glucose uptake was enhanced in cells expression activated STAT5, coinciding with a Hif2 α -dependent upregulation glucose metabolism genes, suggesting that pathways normally active under hypoxia might be utilized by STAT5 under normoxic conditions as well to maintain stem cell properties.

The phenotype imposed on cells by STAT5 might well depend on the actual level of STAT5 activity that is induced. Using a 4-hydroxytamoxifen inducible system that allowed titration of STAT5 activity in human stem and progenitor cells we have demonstrated that the stem cell maintenance properties require intermediate STAT5 activation.¹⁰² On the other hand, high STAT5 activation levels resulted in erythroid differentiation at the expense of HSC self-renewal.¹⁰² C/ebp α levels were reduced upon STAT5 activation, which reached maximum reduction levels at intermediate STAT5 activation. In line with these observations, in mice C/ebp α deficiency resulted in hyperproliferation of hematopoietic progenitor cells and enhancement of hematopoietic stem cell repopulating capacity and self-renewal.^{103,104} Reintroduction of C/ebp α in STAT5A(1*6)-transduced human CD34⁺ cells was sufficient to impair HSC self-renewal capacity.¹⁰⁵ The mechanisms by which STAT5 affects C/ebp α expression levels are still under investigation, but these observations leave open the possibility that enhanced HSC self-renewal might in part be explained by reduction in C/ebp α expression levels.

It is remarkable that the effects of STAT5 on HSC self-renewal are confined to intermediate STAT5 activation levels.¹⁰² This dosage effect of STAT5 on self-renewal is consistent with the observed constitutive activation of STAT5 in AML samples, which is typically lower than cytokine-induced STAT5 activation.^{37,106,107} Such a dosage effect of transcription factors is at present not well understood, but besides STAT5 this has also been observed for the myeloid transcription factor Pu.1, which at 20% expression gives rise to self-renewing murine myeloid leukemias, whereas 50% or 100% reduction in expression do not have such a dramatic effect.^{108,109} Recently, also for Wnt signaling it was demonstrated that intermediate activation levels enhance

self-renewal of HSCs.¹¹⁰ Collectively, these examples clearly underscore the role of transcription factor dosage in regulating HSC self-renewal.

Cell Type-Specific STAT5 Signaling: Differential Role of STAT5 in Hematopoietic Stem and Progenitor Cells?

The observation that STAT5 drives cell cycle progression in various cell types and anti-apoptosis in others, while STAT5 is also required to maintain quiescence of hematopoietic stem cells, suggests that the cell-biological consequences of STAT5 signaling might be highly cell type-specific. We have addressed this issue by introducing a 4-hydroxytamoxifen (4OHT)-inducible STAT5-ER fusion in human stem and progenitor cells.¹⁰¹ Activation of STAT5 specifically in HSC, common myeloid (CMP), granulocyte-macrophage (GMP) or megakaryocyte-erythroid progenitor (MEP) populations resulted in rather distinct phenotypes. Long-term self-renewal and enhanced cobblestone formation could only be imposed on HSCs, but not on committed progenitor subpopulations. Erythroid differentiation could be induced in HSC, CMP and MEP populations, but not in GMPs. Gene

expression profiling revealed that rather distinct gene expression profiles were induced in HSC as compared with more committed progenitor subpopulations. For instance, *Tubb1*, *Hif2α*, *Sod2*, *IL8* and also the cell cycle inhibitor *Cdkn1a/p21* were particularly upregulated in HSCs but not in committed progenitors (Fig. 2). In contrast, *Osm*, *Pim1* and the negative feedback regulators *CISH* and *Socs2* were upregulated both in HSCs and MPPs. The underlying mechanisms are currently unclear, but a number of possibilities might be hypothesized. First, it has been shown that several cofactors such as p300/Cbp, but also interactions with other transcription factors such as Foxo3a, can modulate and fine-tune the STAT5 response.¹¹¹⁻¹¹³ Cell type-specific interaction with such cofactors would then dictate a cell type-specific STAT5 activation pattern of target genes. Seen from this perspective, the modulation of STAT5 signaling by p300/Cbp could add to a stem vs. progenitor-specific component of STAT5. It has been observed that, unlike p300, Cbp is essential for HSC self-renewal maintenance, while p300 is suggested to play a role in differentiation.¹¹⁴⁻¹¹⁶ Interaction of various transcription factors with p300/Cbp is facilitated by the p300/Cbp interacting protein Cited2 which has been shown to be a target gene of STAT5¹¹³ and has differential expression and functions in hematopoietic




Figure 2. Cell type-specific STAT5 signaling in hematopoietic stem and progenitor cells. (1) Expression/activation of cofactors: complex composition. (2) Epigenetic factors that influence STAT5 DNA binding. (3) Expression of receptors and ligands. (4) Niche interactions.

stem vs. progenitor cells¹¹⁷ (and our own observations). In part, such interactions can be mediated by posttranslational modifications such as serine phosphorylation or glycosylation of STAT5.¹¹² Thus, besides the induction of STAT5 tyrosine phosphorylation required for dimerization, nuclear translocation and DNA binding, the simultaneous activation of pathways that mediate STAT5 serine phosphorylation or glycosylation would be required.

Furthermore, the epigenetic status of the cell might play an important role. Hypermethylation of specific promoters or polycomb-mediated condensation of chromatin might prevent STAT5 association with regulatory promoter elements and thus transactivation of certain genes. Clearly, such differences in epigenetic status and cofactor expression might also be dictated by different responses to extracellular stimuli. Thus, the repertoire of specific cytokine and growth factor receptors that is expressed on a cell, as well as direct interactions between hematopoietic stem cells and their bone marrow niche, might ultimately determine the specific STAT5 response (Fig. 2).

Single cell tyrosine phospho-STAT5 analysis revealed that within the normal hematopoietic stem cell and progenitor compartment highly distinct cytokine-induced STAT5 activation patterns are observed.⁶ Also in primary AML patient samples, rather heterogeneous responses toward a series of cytokines were observed, not directly linked to whether or not the cognate receptor was expressed.⁶ There was clear heterogeneity between different patient samples, but also different responses could be observed within distinct cellular compartments within a single patient.^{6,118} For instance, in some patient samples strong IL3 and GM-CSF responses were observed, but only in the CD34⁺ subpopulation, while in other cases strong TPO responses were observed within CD34⁺/CD38⁻ and CD34⁺/CD38⁺ compartments. These observations clearly indicate that strong differences exist in how cytokine and growth factor signals are mediated within a certain cell type, both normal as well as leukemic.

Although elucidation of molecular mechanisms by which cell type specific STAT5 signaling is orchestrated needs further studies, cell type-specific STAT5 target genes clearly do exist. The observation that p21 is upregulated by STAT5, particularly in HSCs, is remarkable (our unpublished observations and ref. 101). It will be interesting to analyze whether the enhanced long-term self-renewal that is observed upon activation of STAT5 in hematopoietic stem cells¹⁶ involves improved stem cell maintenance by keeping the HSCs pool in a relatively quiescent state via upregulation of p21. Knockout studies in mice have indicated that p21 is required during stress hematopoiesis,¹¹⁹ and although p21 was also initially downregulated in STAT5 depleted LSK cells, this downmodulation was not maintained.⁸⁸ On the other hand, in murine embryonic fibroblasts it has also been shown that STAT5 can negatively regulate cell cycle progression through activation of p21.¹²⁰ Inhibition of JAK2/STAT5 signaling by the specific Jak2 inhibitor AZ960 stimulated cell cycling in CD34⁺/CD38⁻ cells in conjunction with downregulation of p21.¹¹⁸ Further, activation of p21 has been shown to be critical in preventing excess DNA-damage accumulation and functional exhaustion of leukemic stem cells,¹²¹ and it will be interesting to further reveal its role downstream of STAT5 in HSCs.

Furthermore, HIF2 α was upregulated in HSCs and CMPs by STAT5, but not in MEPs and GMPs.¹⁰¹ Under normoxic conditions, proline residues of Hypoxia-Induced Factor 2 are hydroxylated resulting in a reduction in protein levels via VHL-mediated proteasomal degradation. Under hypoxic conditions, such as in the presumed endosteal quiescent stem cell niche, Hifs are stabilized and act as transcription factors.¹²² It is currently unknown whether and which Hif-induced target genes are essential to maintain stemness of normal HSCs, but it was recently shown that in Hif1^{-/-} mice HSCs numbers decrease during stress which was associated with a loss of HSC quiescence.¹²³ Another report indicated that HSCs in the quiescence niche utilize glycolysis for their energy demands, which depended on a Meis1-induced Hif1 α signaling network.¹²⁴ Whether Hif1 α and Hif2 α display similar or distinct functions in HSCs remains to be established.

Our understanding of the mechanisms that determine whether, where and when a stem cell will self-renew or differentiate is still limited, but recent advances have indicated that the stem cell microenvironment provides essential cues that direct these cell fate decisions.¹²⁵⁻¹²⁸ It is remarkable that STAT5-induced long-term self-renewal is typically observed when cells are cultured in direct contact with stromal cells,¹⁶ in contrast to e.g., Bmi1-induced self-renewal which occurred in a more microenvironment-independent manner.¹²⁹ Thus, altered interactions with the stem cells niche might also underlie the enhanced self-renewal properties imposed on HSCs by activated STAT5. Although the mechanisms by which the interaction with the microenvironment of STAT5A(1*6)-expressing CD34⁺ cells are still unclear, our ongoing studies in which gene-expression profiling was performed in HSCs and progenitor subsets revealed that the list of STAT5-targets is significantly enriched for membrane (-associated) proteins.¹⁰² One of the STAT5 targets that has been identified is MUC1¹³⁰ which is a (proto)oncogene involved in adhesion and transendothelial migration, and has been associated with initiation of various intracellular signal transduction pathways including β -Catenin, p53 and NF κ B pathways.¹³¹⁻¹³⁶ Also, MUC1 has been shown to mediate an oscillatory calcium signal upon binding to ICAM1.¹³⁷ Within the endosteal region of the bone marrow where stem cells are thought to reside, Ca²⁺ levels are high, and HSC retention within the niche depends on the Calcium-Sensing receptor (CaR).¹³⁸ Thus, STAT5 might exert its phenotype, at least in part, by influencing interactions between HSCs and their niche.

Conclusions and Future Perspectives

In both murine and human model systems it has been convincingly shown that STAT5 fulfills an important role in hematopoietic stem cell self-renewal. Although the precise mechanisms by which HSC self-renewal is orchestrated by STAT5 remain elusive till date, an increasing number of STAT5 target genes have been identified that are currently under investigation. In myeloproliferative diseases and leukemias, a number of oncogenes have been identified that are capable of inducing STAT5 activity, and accumulating evidence has

indicated that STAT5 participates in self-renewal of leukemic stem cells as well. Thus, it appears likely that STAT5 will become an important diagnostic marker in the near future, and

specific targeting of STAT5 should be focus of therapeutical intervention strategies to improve treatment of hematological malignancies.

References

- Bunting KD. STAT5 signaling in normal and pathologic hematopoiesis. *Front Biosci* 2007; 12:2807-20; PMID: 17485261; <http://dx.doi.org/10.2741/2274>
- Paukku K, Silvennoinen O. STATs as critical mediators of signal transduction and transcription: lessons learned from STAT5. *Cytokine Growth Factor Rev* 2004; 15:435-55; PMID:15561601; <http://dx.doi.org/10.1016/j.cytogfr.2004.09.001>
- Schindler C, Darnell JE, Jr. Transcriptional responses to polypeptide ligands: the JAK-STAT pathway. *Annu Rev Biochem* 1995; 64:621-52; PMID:7574495; <http://dx.doi.org/10.1146/annurev.bi.64.070195.003201>
- Ihle JN. The Stat family in cytokine signaling. *Curr Opin Cell Biol* 2001; 13:211-7; PMID:11248555; [http://dx.doi.org/10.1016/S0955-0674\(00\)00199-X](http://dx.doi.org/10.1016/S0955-0674(00)00199-X)
- Hennighausen L, Robinson GW. Interpretation of cytokine signaling through the transcription factors STAT5A and STAT5B. *Genes Dev* 2008; 22:711-21; PMID: 18347089; <http://dx.doi.org/10.1101/gad.1643908>
- Han L, Wierenga AT, Rozenveld-Geugien M, van de Lande K, Vellenga E, Schuringa JJ. Single-cell STAT5 signal transduction profiling in normal and leukemic stem and progenitor cell populations reveals highly distinct cytokine responses. *PLoS One* 2009; 4:e7989; PMID:19956772; <http://dx.doi.org/10.1371/journal.pone.0007989>
- Ryan JJ, Huang H, McReynolds LJ, Shelburne C, Hu-Li J, Huff TF, et al. Stem cell factor activates STAT-5 DNA binding in IL-3-derived bone marrow mast cells. *Exp Hematol* 1997; 25:357-62; PMID: 9131012
- Pallard C, Gouilleux F, B nit L, Cocault L, Souyri M, Levy D, et al. Thrombopoietin activates a STAT5-like factor in hematopoietic cells. *EMBO J* 1995; 14:2847-56; PMID:7796811
- Feugier P, Li N, Jo DY, Shieh JH, MacKenzie KL, Lesesve JF, et al. Osteopetrotic mouse stroma with thrombopoietin, c-kit ligand, and flk-2 ligand supports long-term mobilized CD34+ hematopoiesis in vitro. *Stem Cells Dev* 2005; 14:505-16; PMID:16305336; <http://dx.doi.org/10.1089/scd.2005.14.505>
- Seita J, Ema H, Oechara J, Yamazaki S, Tadokoro Y, Yamasaki A, et al. Lnk negatively regulates self-renewal of hematopoietic stem cells by modifying thrombopoietin-mediated signal transduction. *Proc Natl Acad Sci U S A* 2007; 104:2349-54; PMID:17284614; <http://dx.doi.org/10.1073/pnas.0606238104>
- Pallard C, Gouilleux F, Charon M, Groner B, Gisselbrecht S, Dusanter-Fourt I. Interleukin-3, erythropoietin, and prolactin activate a STAT5-like factor in lymphoid cells. *J Biol Chem* 1995; 270:15942-5; PMID:7608147; <http://dx.doi.org/10.1074/jbc.270.27.15942>
- Nosaka T, Kawashima T, Misawa K, Ikuta K, Mui AL, Kitamura T. STAT5 as a molecular regulator of proliferation, differentiation and apoptosis in hematopoietic cells. *EMBO J* 1999; 18:4754-65; PMID: 10469654; <http://dx.doi.org/10.1093/emboj/18.17.4754>
- Socolovsky M, Fallon AE, Wang S, Brugnara C, Lodish HF. Fetal anemia and apoptosis of red cell progenitors in Stat5a^{-/-} Stat5b^{-/-} mice: a direct role for Stat5 in Bcl-X(L) induction. *Cell* 1999; 98:181-91; PMID:10428030; [http://dx.doi.org/10.1016/S0092-8674\(00\)81013-2](http://dx.doi.org/10.1016/S0092-8674(00)81013-2)
- Socolovsky M, Nam H, Fleming MD, Haase VH, Brugnara C, Lodish HF. Ineffective erythropoiesis in Stat5a^{-/-} Stat5b^{-/-} mice due to decreased survival of early erythroblasts. *Blood* 2001; 98:3261-73; PMID: 11719363; <http://dx.doi.org/10.1182/blood.V98.12.3261>
- Gar on L, Rivat C, James C, Lacout C, Camara-Clayette V, Ugo V, et al. Constitutive activation of STAT5 and Bcl-xL overexpression can induce endogenous erythroid colony formation in human primary cells. *Blood* 2006; 108:1551-4; PMID:16684963; <http://dx.doi.org/10.1182/blood-2005-10-009514>
- Schuringa JJ, Chung KY, Morrone G, Moore MA. Constitutive activation of STAT5A promotes human hematopoietic stem cell self-renewal and erythroid differentiation. *J Exp Med* 2004; 200:623-35; PMID: 15353555; <http://dx.doi.org/10.1084/jem.20041024>
- Schuringa JJ, Wu K, Morrone G, Moore MA. Enforced activation of STAT5A facilitates the generation of embryonic stem-derived hematopoietic stem cells that contribute to hematopoiesis in vivo. *Stem Cells* 2004; 22:1191-204; PMID:15579639; <http://dx.doi.org/10.1634/stemcells.2004-0033>
- Grebien F, Kerenyi MA, Kovacic B, Kolbe T, Becker V, Dolznig H, et al. Stat5 activation enables erythropoiesis in the absence of EpoR and Jak2. *Blood* 2008; 111:4511-22; PMID:18239084; <http://dx.doi.org/10.1182/blood-2007-07-102848>
- Li G, Wang Z, Zhang Y, Kang Z, Haviernikova E, Cui Y, et al. STAT5 requires the N-domain to maintain hematopoietic stem cell repopulating function and appropriate lymphoid-myeloid lineage output. *Exp Hematol* 2007; 35:1684-94; PMID:17976521; <http://dx.doi.org/10.1016/j.exphem.2007.08.026>
- Mui AL, Wakao H, O'Farrell AM, Harada N, Miyajima A. Interleukin-3, granulocyte-macrophage colony stimulating factor and interleukin-5 transduce signals through two STAT5 homologs. *EMBO J* 1995; 14:1166-75; PMID:7720707
- Azam M, Erdjument-Bromage H, Kreider BL, Xia M, Quelle F, Basu R, et al. Interleukin-3 signals through multiple isoforms of Stat5. *EMBO J* 1995; 14:1402-11; PMID:7537213
- Coffer PJ, Koenderman L, de Groot RP. The role of STATs in myeloid differentiation and leukemia. *Oncogene* 2000; 19:2511-22; PMID:10851050; <http://dx.doi.org/10.1038/sj.onc.1203479>
- Kieslinger M, Woldman I, Moriggl R, Hofmann J, Marine JC, Ihle JN, et al. Antiapoptotic activity of Stat5 required during terminal stages of myeloid differentiation. *Genes Dev* 2000; 14:232-44; PMID:10652277
- Xiao W, Hong H, Kawakami Y, Lowell CA, Kawakami T. Regulation of myeloproliferation and M2 macrophage programming in mice by Lyn/Hck, SHIP, and Stat5. *J Clin Invest* 2008; 118:924-34; PMID:18246197
- Kimura A, Rieger MA, Simone JM, Chen W, Wickre MC, Zhu BM, et al. The transcription factors STAT5A/B regulate GM-CSF-mediated granulopoiesis. *Blood* 2009; 114:4721-8; PMID:19779039; <http://dx.doi.org/10.1182/blood-2009-04-216390>
- Buitenhuis M, Baltus B, Lammers JW, Coffer PJ, Koenderman L. Signal transducer and activator of transcription 5a (STAT5a) is required for eosinophil differentiation of human cord blood-derived CD34+ cells. *Blood* 2003; 101:134-42; PMID:12393707; <http://dx.doi.org/10.1182/blood-2002-03-0740>
- Hoelbl A, Kovacic B, Kerenyi MA, Simma O, Warsch W, Cui Y, et al. Clarifying the role of Stat5 in lymphoid development and Abelson-induced transformation. *Blood* 2006; 107:4898-906; PMID:16493008; <http://dx.doi.org/10.1182/blood-2005-09-3596>
- Moriggl R, Sexl V, Piekorz R, Topham D, Ihle JN. Stat5 activation is uniquely associated with cytokine signaling in peripheral T cells. *Immunity* 1999; 11:225-30; PMID:10485657; [http://dx.doi.org/10.1016/S1074-7613\(00\)80097-7](http://dx.doi.org/10.1016/S1074-7613(00)80097-7)
- Moriggl R, Topham DJ, Teglund S, Sexl V, McKay C, Wang D, et al. Stat5 is required for IL-2-induced cell cycle progression of peripheral T cells. *Immunity* 1999; 10:249-59; PMID:10072077; [http://dx.doi.org/10.1016/S1074-7613\(00\)80025-4](http://dx.doi.org/10.1016/S1074-7613(00)80025-4)
- Dai X, Chen Y, Di L, Podd A, Li G, Bunting KD, et al. Stat5 is essential for early B cell development but not for B cell maturation and function. *J Immunol* 2007; 179:1068-79; PMID:17617599
- Rocnik JL, Okabe R, Yu JC, Lee BH, Giese N, Schenkein DP, et al. Roles of tyrosine 589 and 591 in STAT5 activation and transformation mediated by FLT3-ITD. *Blood* 2006; 108:1339-45; PMID:16627759; <http://dx.doi.org/10.1182/blood-2005-11-011429>
- Chung KY, Morrone G, Schuringa JJ, Wong B, Dorn DC, Moore MA. Enforced expression of an Flt3 internal tandem duplication in human CD34+ cells confers properties of self-renewal and enhanced erythropoiesis. *Blood* 2005; 105:77-84; PMID:15242879; <http://dx.doi.org/10.1182/blood-2003-12-4445>
- Schepers H, van Goslina D, Wierenga AT, Eggen BJ, Schuringa JJ, Vellenga E. STAT5 is required for long-term maintenance of normal and leukemic human stem/progenitor cells. *Blood* 2007; 110:2880-8; PMID: 17630355; <http://dx.doi.org/10.1182/blood-2006-08-039073>
- Choudhary C, Brandts C, Schwable J, Tickenbrock L, Sargin B, Ueker A, et al. Activation mechanisms of STAT5 by oncogenic Flt3-ITD. *Blood* 2007; 110:370-4; PMID:17356133; <http://dx.doi.org/10.1182/blood-2006-05-024018>
- Choudhary C, Schwable J, Brandts C, Tickenbrock L, Sargin B, Kindler T, et al. AML-associated Flt3 kinase domain mutations show signal transduction differences compared with Flt3 ITD mutations. *Blood* 2005; 106:265-73; PMID:15769897; <http://dx.doi.org/10.1182/blood-2004-07-2942>
- Kiyoi H, Ohno R, Ueda R, Saito H, Naoe T. Mechanism of constitutive activation of FLT3 with internal tandem duplication in the juxtamembrane domain. *Oncogene* 2002; 21:2555-63; PMID: 11971190; <http://dx.doi.org/10.1038/sj.onc.1205332>
- Birkenkamp KU, Geugien M, Lemmink HH, Kruijer W, Vellenga E. Regulation of constitutive STAT5 phosphorylation in acute myeloid leukemia blasts. *Leukemia* 2001; 15:1923-31; PMID:11753614; <http://dx.doi.org/10.1038/sj.leu.2402317>
- Mizuki M, Fenski R, Halfter H, Matsumura I, Schmidt R, M ller C, et al. Flt3 mutations from patients with acute myeloid leukemia induce transformation of 32D cells mediated by the Ras and STAT5 pathways. *Blood* 2000; 96:3907-14; PMID: 11090077
- Abu-Duhier FM, Goodeve AC, Wilson GA, Care RS, Peake IR, Reilly JT. Identification of novel FLT-3 Asp835 mutations in adult acute myeloid leukaemia. *Br J Haematol* 2001; 113:983-8; PMID:11442493; <http://dx.doi.org/10.1046/j.1365-2141.2001.02850.x>
- Schwaller J, Parganas E, Wang D, Cain D, Aster JC, Williams IR, et al. Stat5 is essential for the myelo- and lymphoproliferative disease induced by TEL/JAK2. *Mol Cell* 2000; 6:693-704; PMID:11030348; [http://dx.doi.org/10.1016/S1097-2765\(00\)00067-8](http://dx.doi.org/10.1016/S1097-2765(00)00067-8)
- Baxter EJ, Scott LM, Campbell PJ, East C, Fourouclas N, Swanton S, et al. Cancer Genome Project. Acquired mutation of the tyrosine kinase JAK2 in human myeloproliferative disorders. *Lancet* 2005; 365:1054-61; PMID:15781101

42. James C, Ugo V, Le Couédic JP, Staerk J, Delhommeau F, Lacout C, et al. A unique clonal JAK2 mutation leading to constitutive signalling causes polycythemia vera. *Nature* 2005; 434:1144-8; PMID: 15793561; <http://dx.doi.org/10.1038/nature03546>
43. Kralovics R, Passamonti F, Buser AS, Teo SS, Tiedt R, Passweg JR, et al. A gain-of-function mutation of JAK2 in myeloproliferative disorders. *N Engl J Med* 2005; 352:1779-90; PMID:15858187; <http://dx.doi.org/10.1056/NEJMoa051113>
44. Levine RL, Wadleigh M, Cools J, Ebert BL, Wernig G, Huntly BJ, et al. Activating mutation in the tyrosine kinase JAK2 in polycythemia vera, essential thrombocythemia, and myeloid metaplasia with myelofibrosis. *Cancer Cell* 2005; 7:387-97; PMID:15837627; <http://dx.doi.org/10.1016/j.ccr.2005.03.023>
45. Walters DK, Mercher T, Gu TL, O'Hare T, Tyner JW, Loriaux M, et al. Activating alleles of JAK3 in acute megakaryoblastic leukemia. *Cancer Cell* 2006; 10:65-75; PMID:16843266; <http://dx.doi.org/10.1016/j.ccr.2006.06.002>
46. Brizzi MF, Dentelli P, Rosso A, Yarden Y, Pegoraro L. STAT protein recruitment and activation in c-Kit deletion mutants. *J Biol Chem* 1999; 274:16965-72; PMID:10358045; <http://dx.doi.org/10.1074/jbc.274.24.16965>
47. Ikeda H, Kanakura Y, Tamaki T, Kuriu A, Kitayama H, Ishikawa J, et al. Expression and functional role of the proto-oncogene c-kit in acute myeloblastic leukemia cells. *Blood* 1991; 78:2962-8; PMID:1720040
48. Beghini A, Larizza L, Cairoli R, Morra E. c-kit activating mutations and mast cell proliferation in human leukemia. *Blood* 1998; 92:701-2; PMID:9657776
49. Corbacioglu S, Kilic M, Westhoff MA, Reinhardt D, Fulda S, Debatin KM. Newly identified c-KIT receptor tyrosine kinase ITD in childhood AML induces ligand-independent growth and is responsive to a synergistic effect of imatinib and rapamycin. *Blood* 2006; 108:3504-13; PMID:16840725; <http://dx.doi.org/10.1182/blood-2006-05-021691>
50. Cain JA, Xiang Z, O'Neal J, Kreisel F, Colson A, Luo H, et al. Myeloproliferative disease induced by TEL-PDGFRB displays dynamic range sensitivity to Stat5 gene dosage. *Blood* 2007; 109:3906-14; PMID: 17218386; <http://dx.doi.org/10.1182/blood-2006-07-036335>
51. Sternberg DW, Tomasson MH, Carroll M, Curley DP, Barker G, Caprio M, et al. The TEL/PDGFRbeta fusion in chronic myelomonocytic leukemia signals through STAT5-dependent and STAT5-independent pathways. *Blood* 2001; 98:3390-7; PMID:11719379; <http://dx.doi.org/10.1182/blood.V98.12.3390>
52. Buitenhuis M, Verhagen LP, Cools J, Coffey PJ. Molecular mechanisms underlying FIP1L1-PDGFR-mediated myeloproliferation. *Cancer Res* 2007; 67: 3759-66; PMID:17440089; <http://dx.doi.org/10.1158/0008-5472.CAN-06-4183>
53. Cools J, Stover EH, Gilliland DG. Detection of the FIP1L1-PDGFR fusion in idiopathic hypereosinophilic syndrome and chronic eosinophilic leukemia. *Methods Mol Med* 2006; 125:177-87; PMID: 16502585
54. Danial NN, Pernis A, Rothman PB. Jak-STAT signaling induced by the v-abl oncogene. *Science* 1995; 269:1875-7; PMID:7569929; <http://dx.doi.org/10.1126/science.7569929>
55. Ye D, Wolff N, Li L, Zhang S, Ilaria RL, Jr. STAT5 signaling is required for the efficient induction and maintenance of CML in mice. *Blood* 2006; 107: 4917-25; PMID:16522816; <http://dx.doi.org/10.1182/blood-2005-10-4110>
56. Heath C, Cross NC. Critical role of STAT5 activation in transformation mediated by ZNF198-FGFR1. *J Biol Chem* 2004; 279:6666-73; PMID:14660670; <http://dx.doi.org/10.1074/jbc.M308743200>
57. Dong S, Twardy DJ. Interactions of STAT5b-RARalpha, a novel acute promyelocytic leukemia fusion protein, with retinoic acid receptor and STAT3 signaling pathways. *Blood* 2002; 99:2637-46; PMID:11929748; <http://dx.doi.org/10.1182/blood.V99.8.2637>
58. Arnould C, Philippe C, Bourdon V, Grigoire MJ, Berger R, Jonveaux P. The signal transducer and activator of transcription STAT5b gene is a new partner of retinoic acid receptor alpha in acute promyelocytic-like leukaemia. *Hum Mol Genet* 1999; 8:1741-9; PMID: 10441338; <http://dx.doi.org/10.1093/hmg/8.9.1741>
59. Carlesso N, Frank DA, Griffin JD. Tyrosyl phosphorylation and DNA binding activity of signal transducers and activators of transcription (STAT) proteins in hematopoietic cell lines transformed by Bcr/Abl. *J Exp Med* 1996; 183:811-20; PMID: 8642285; <http://dx.doi.org/10.1084/jem.183.3.811>
60. Ilaria RL, Jr., Van Etten RA. P210 and P190(BCR/ABL) induce the tyrosine phosphorylation and DNA binding activity of multiple specific STAT family members. *J Biol Chem* 1996; 271:31704-10; PMID: 8940193; <http://dx.doi.org/10.1074/jbc.271.49.31704>
61. Frank DA, Varticovski L. BCR/abl leads to the constitutive activation of Stat proteins, and shares an epitope with tyrosine phosphorylated Stats. *Leukemia* 1996; 10:1724-30; PMID:8892675
62. Shuai K, Halpern J, ten Hoeve J, Rao X, Sawyers CL. Constitutive activation of STAT5 by the BCR-ABL oncogene in chronic myelogenous leukemia. *Oncogene* 1996; 13:247-54; PMID:8710363
63. Nieborowska-Skorska M, Wasik MA, Slupianek A, Salomoni P, Kitamura T, Calabretta B, et al. Signal transducer and activator of transcription (STAT)5 activation by BCR/ABL is dependent on intact Src homology (SH)3 and SH2 domains of BCR/ABL and is required for leukemogenesis. *J Exp Med* 1999; 189:1229-42; PMID:10209040; <http://dx.doi.org/10.1084/jem.189.8.1229>
64. Horita M, Andreu EJ, Benito A, Arbona C, Sanz C, Benet I, et al. Blockade of the Bcr-Abl kinase activity induces apoptosis of chronic myelogenous leukemia cells by suppressing signal transducer and activator of transcription 5-dependent expression of Bcl-xL. *J Exp Med* 2000; 191:977-84; PMID:10727459; <http://dx.doi.org/10.1084/jem.191.6.977>
65. de Groot RP, Raaijmakers JA, Lammers JW, Jove R, Koenderman L. STAT5 activation by BCR-Abl contributes to transformation of K562 leukemia cells. *Blood* 1999; 94:1108-12; PMID:10419904
66. Sillaber C, Gesbert F, Frank DA, Sattler M, Griffin JD. STAT5 activation contributes to growth and viability in Bcr/Abl-transformed cells. *Blood* 2000; 95:2118-25; PMID:10706883
67. Futami M, Harano T, Soda Y, Kobayashi S, Miyagishi M, Tojo A. RNAi-mediated silencing of p190Bcr-Abl inactivates Stat5 and cooperates with imatinib mesylate and 17-allylamino-17-demethoxygeldanamycin in selective killing of p190Bcr-Abl-expressing leukemia cells. *Leukemia* 2008; 22:1131-8; PMID:18368071; <http://dx.doi.org/10.1038/leu.2008.60>
68. Scherr M, Chaturvedi A, Battmer K, Dallmann I, Schultheis B, Ganser A, et al. Enhanced sensitivity to inhibition of SHP2, STAT5, and Gab2 expression in chronic myeloid leukemia (CML). *Blood* 2006; 107:3279-87; PMID:16278304; <http://dx.doi.org/10.1182/blood-2005-08-3087>
69. Warsch W, Kollmann K, Eckelhart E, Fajmann S, Cerny-Reiterer S, Höbl A, et al. High STAT5 levels mediate imatinib resistance and indicate disease progression in chronic myeloid leukemia. *Blood* 2011; 117:3409-20; PMID:21220747; <http://dx.doi.org/10.1182/blood-2009-10-248211>
70. Hoelbl A, Schuster C, Kovacic B, Zhu B, Wickre M, Hoelzl MA, et al. Stat5 is indispensable for the maintenance of bcr/abl-positive leukaemia. *EMBO Mol Med* 2010; 2:98-110; PMID:20201032; <http://dx.doi.org/10.1002/emmm.201000062>
71. Nelson EA, Walker SR, Weisberg E, Bar-Natan M, Barrett R, Gashin LB, et al. The STAT5 inhibitor pimozone decreases survival of chronic myelogenous leukemia cells resistant to kinase inhibitors. *Blood* 2011; 117:3421-9; PMID:21233313; <http://dx.doi.org/10.1182/blood-2009-11-255232>
72. Aboudola S, Murugesan G, Szpurka H, Ramsingh G, Zhao X, Prescott N, et al. Bone marrow phospho-STAT5 expression in non-CML chronic myeloproliferative disorders correlates with JAK2 V617F mutation and provides evidence of in vivo JAK2 activation. *Am J Surg Pathol* 2007; 31:233-9; PMID:17255768; <http://dx.doi.org/10.1097/01.pas.0000213338.25111.d3>
73. Wernig G, Kharas MG, Okabe R, Moore SA, Leeman DS, Cullen DE, et al. Efficacy of TG101348, a selective JAK2 inhibitor, in treatment of a murine model of JAK2V617F-induced polycythemia vera. *Cancer Cell* 2008; 13:311-20; PMID:18394554; <http://dx.doi.org/10.1016/j.ccr.2008.02.009>
74. Geron I, Abrahamson AE, Barroga CF, Kavalchik E, Gotlib J, Hood JD, et al. Selective inhibition of JAK2-driven erythroid differentiation of polycythemia vera progenitors. *Cancer Cell* 2008; 13:321-30; PMID: 18394555; <http://dx.doi.org/10.1016/j.ccr.2008.02.017>
75. Moriggl R, Sexl V, Kenner L, Dunsch C, Stangl K, Gingras S, et al. Stat5 tetramer formation is associated with leukemogenesis. *Cancer Cell* 2005; 7:87-99; PMID: 15652752; <http://dx.doi.org/10.1016/j.ccr.2004.12.010>
76. Onishi M, Nosaka T, Misawa K, Mui AL, Gorman D, McMahon M, et al. Identification and characterization of a constitutively active STAT5 mutant that promotes cell proliferation. *Mol Cell Biol* 1998; 18:3871-9; PMID:9632771
77. Kato Y, Iwama A, Tadokoro Y, Shimoda K, Minoguchi M, Akira S, et al. Selective activation of STAT5 unveils its role in stem cell self-renewal in normal and leukemic hematopoiesis. *J Exp Med* 2005; 202:169-79; PMID:15998795; <http://dx.doi.org/10.1084/jem.20042541>
78. Chalandon Y, Jiang X, Christ O, Loutet S, Thanopoulou E, Eaves A, et al. BCR-ABL-transduced human cord blood cells produce abnormal populations in immunodeficient mice. *Leukemia* 2005; 19:442-8; PMID:15674417; <http://dx.doi.org/10.1038/sj.leu.2403650>
79. Rizo A, Horton SJ, Olthoff S, Dontje B, Aulsebrook A, van Os R, et al. BM1 collaborates with BCR-ABL in leukemic transformation of human CD34+ cells. *Blood* 2010; 116:4621-30; PMID:20724541; <http://dx.doi.org/10.1182/blood-2010-02-270660>
80. Bunting KD, Bradley HL, Hawley TS, Moriggl R, Sorrentino BP, Ihle JN. Reduced lymphomyeloid repopulating activity from adult bone marrow and fetal liver of mice lacking expression of STAT5. *Blood* 2002; 99:479-87; PMID:11781228; <http://dx.doi.org/10.1182/blood.V99.2.479>
81. Bradley HL, Hawley TS, Bunting KD. Cell intrinsic defects in cytokine responsiveness of STAT5-deficient hematopoietic stem cells. *Blood* 2002; 100:3983-9; PMID:12393407; <http://dx.doi.org/10.1182/blood-2002-05-1602>
82. Bradley HL, Couldrey C, Bunting KD. Hematopoietic-repopulating defects from STAT5-deficient bone marrow are not fully accounted for by loss of thrombopoietin responsiveness. *Blood* 2004; 103:2965-72; PMID:15070672; <http://dx.doi.org/10.1182/blood-2003-08-2963>

83. Snow JW, Abraham N, Ma MC, Abbey NW, Herndier B, Goldsmith MA. STAT5 promotes multi-lineage hematolymphoid development in vivo through effects on early hematopoietic progenitor cells. *Blood* 2002; 99:95-101; PMID:11756158; <http://dx.doi.org/10.1182/blood.V99.1.95>
84. Snow JW, Abraham N, Ma MC, Bronson SK, Goldsmith MA. Transgenic bcl-2 is not sufficient to rescue all hematolymphoid defects in STAT5A/5B-deficient mice. *Exp Hematol* 2003; 31:1253-8; PMID:14662332; <http://dx.doi.org/10.1016/j.exphem.2003.09.014>
85. Snow JW, Abraham N, Ma MC, Goldsmith MA. Bone marrow transplant completely rescues hematolymphoid defects in STAT5A/5B-deficient mice. *Exp Hematol* 2003; 31:1247-52; PMID:14662331; <http://dx.doi.org/10.1016/j.exphem.2003.08.013>
86. Couldrey C, Bradley HL, Bunting KDA. A STAT5 modifier locus on murine chromosome 7 modulates engraftment of hematopoietic stem cells during steady-state hematopoiesis. *Blood* 2005; 105:1476-83; PMID:15498858; <http://dx.doi.org/10.1182/blood-2004-06-2302>
87. Zhu BM, McLaughlin SK, Na R, Liu J, Cui Y, Martin C, et al. Hematopoietic-specific Stat5-null mice display microcytic hypochromic anemia associated with reduced transferrin receptor gene expression. *Blood* 2008; 112:2071-80; PMID:18552213; <http://dx.doi.org/10.1182/blood-2007-12-127480>
88. Wang Z, Li G, Tse W, Bunting KD. Conditional deletion of STAT5 in adult mouse hematopoietic stem cells causes loss of quiescence and permits efficient nonablative stem cell replacement. *Blood* 2009; 113:4856-65; PMID:19258595; <http://dx.doi.org/10.1182/blood-2008-09-181107>
89. Matikainen S, Sareneva T, Ronni T, Lehtonen A, Koskinen PJ, Julkunen I. Interferon-alpha activates multiple STAT proteins and upregulates proliferation-associated IL-2Ralpha, c-myc, and pim-1 genes in human T cells. *Blood* 1999; 93:1980-91; PMID:10068671
90. Matsumura I, Kitamura T, Wakao H, Tanaka H, Hashimoto K, Albanese C, et al. Transcriptional regulation of the cyclin D1 promoter by STAT5: its involvement in cytokine-dependent growth of hematopoietic cells. *EMBO J* 1999; 18:1367-77; PMID:10064602; <http://dx.doi.org/10.1093/emboj/18.5.1367>
91. Yip-Schneider MT, Horie M, Broxmeyer HE. Transcriptional induction of pim-1 protein kinase gene expression by interferon gamma and post-transcriptional effects on costimulation with steel factor. *Blood* 1995; 85:3494-502; PMID:7540064
92. Zozar K, Ciemerych MA, Rebel VI, Shigematsu H, Zagodzón A, Sicinska E, et al. Mouse development and cell proliferation in the absence of D-cyclins. *Cell* 2004; 118:477-91; PMID:15315760; <http://dx.doi.org/10.1016/j.cell.2004.07.025>
93. Milkers H, Nawijn M, Allen J, Brouwers C, Verhoeven E, Jonkers J, et al. Mice deficient for all PIM kinases display reduced body size and impaired responses to hematopoietic growth factors. *Mol Cell Biol* 2004; 24:6104-15; PMID:15199164; <http://dx.doi.org/10.1128/MCB.24.13.6104-6115.2004>
94. Mochizuki T, Kitanaka K, Noguchi K, Muramatsu T, Asai A, Kuchino Y. Physical and functional interactions between Pim-1 kinase and Cdc25A phosphatase. Implications for the Pim-1-mediated activation of the c-Myc signaling pathway. *J Biol Chem* 1999; 274:18659-66; PMID:10373478; <http://dx.doi.org/10.1074/jbc.274.26.18659>
95. Bachmann M, Kosan C, Xing PX, Montenarh M, Hoffmann I, Möry T. The oncogenic serine/threonine kinase Pim-1 directly phosphorylates and activates the G2/M specific phosphatase Cdc25C. *Int J Biochem Cell Biol* 2006; 38:430-43; PMID:16356754; <http://dx.doi.org/10.1016/j.biocel.2005.10.010>
96. Satoh Y, Matsumura I, Tanaka H, Ezoe S, Sugahara H, Mizuki M, et al. Roles for c-Myc in self-renewal of hematopoietic stem cells. *J Biol Chem* 2004; 279:24986-93; PMID:15067010; <http://dx.doi.org/10.1074/jbc.M400407200>
97. Wilson A, Murphy MJ, Oskarsson T, Kaloulis K, Bettess MD, Oser GM, et al. c-Myc controls the balance between hematopoietic stem cell self-renewal and differentiation. *Genes Dev* 2004; 18:2747-63; PMID:15545632; <http://dx.doi.org/10.1101/gad.313104>
98. Yoshimoto G, Miyamoto T, Jabbarzadeh-Tabrizi S, Iino T, Rocnik JL, Kikushige Y, et al. FLT3-ITD up-regulates MCL-1 to promote survival of stem cells in acute myeloid leukemia via FLT3-ITD-specific STAT5 activation. *Blood* 2009; 114:5034-43; PMID:19808698; <http://dx.doi.org/10.1182/blood-2008-12-196055>
99. Xu M, Nie L, Kim SH, Sun XH. STAT5-induced Id-1 transcription involves recruitment of HDAC1 and deacetylation of C/EBPbeta. *EMBO J* 2003; 22:893-904; PMID:12574125; <http://dx.doi.org/10.1093/emboj/cdg094>
100. Jankovic V, Ciarrocchi A, Boccini P, DeBlasio T, Benezra R, Nimer SD. Id1 restrains myeloid commitment, maintaining the self-renewal capacity of hematopoietic stem cells. *Proc Natl Acad Sci U S A* 2007; 104:1260-5; PMID:17227850; <http://dx.doi.org/10.1073/pnas.0607894104>
101. Fatrai S, Wierenga AT, Daenen SM, Vellenga E, Schuringa JJ. Identification of HIF2alpha as an important STAT5 target gene in human hematopoietic stem cells. *Blood* 2011; 117:3320-30; PMID:21263150; <http://dx.doi.org/10.1182/blood-2010-08-303669>
102. Wierenga AT, Vellenga E, Schuringa JJ. Maximal STAT5-induced proliferation and self-renewal at intermediate STAT5 activity levels. *Mol Cell Biol* 2008; 28:6668-80; PMID:18779318; <http://dx.doi.org/10.1128/MCB.01025-08>
103. Heath V, Suh HC, Holman M, Renn K, Gooya JM, Parkin S, et al. C/EBPalpha deficiency results in hyperproliferation of hematopoietic progenitor cells and disrupts macrophage development in vitro and in vivo. *Blood* 2004; 104:1639-47; PMID:15073037; <http://dx.doi.org/10.1182/blood-2003-11-3963>
104. Zhang P, Iwasaki-Arai J, Iwasaki H, Fenyus ML, Dayaram T, Owens BM, et al. Enhancement of hematopoietic stem cell repopulating capacity and self-renewal in the absence of the transcription factor C/EBP alpha. *Immunity* 2004; 21:853-63; PMID:15589173; <http://dx.doi.org/10.1016/j.immuni.2004.11.006>
105. Schepers H, Wierenga AT, van Gosliga D, Eggen BJ, Vellenga E, Schuringa JJ. Reintroduction of C/EBPalpha in leukemic CD34+ stem/progenitor cells impairs self-renewal and partially restores myelopoiesis. *Blood* 2007; 110:1317-25; PMID:17475913; <http://dx.doi.org/10.1182/blood-2006-10-052175>
106. Spiekermann K, Pau M, Schwab R, Schmiejia K, Franzrahe S, Hiddemann W. Constitutive activation of STAT3 and STAT5 is induced by leukemic fusion proteins with protein tyrosine kinase activity and is sufficient for transformation of hematopoietic precursor cells. *Exp Hematol* 2002; 30:262-71; PMID:11882364; [http://dx.doi.org/10.1016/S0301-472X\(01\)00787-1](http://dx.doi.org/10.1016/S0301-472X(01)00787-1)
107. Harir N, Pecquet C, Kerenyi M, Sonneck K, Kovacic B, Nyrga R, et al. Constitutive activation of Stat5 promotes its cytoplasmic localization and association with PI3-kinase in myeloid leukemias. *Blood* 2007; 109:1678-86; PMID:17038539; <http://dx.doi.org/10.1182/blood-2006-01-029918>
108. Rosenbauer F, Wagner K, Kutok JL, Iwasaki H, Le Beau MM, Okuno Y, et al. Acute myeloid leukemia induced by graded reduction of a lineage-specific transcription factor, PU.1. *Nat Genet* 2004; 36:624-30; PMID:15146183; <http://dx.doi.org/10.1038/ng1361>
109. Rosenbauer F, Owens BM, Yu L, Tumang JR, Steidl U, Kutok JL, et al. Lymphoid cell growth and transformation are suppressed by a key regulatory element of the gene encoding PU.1. *Nat Genet* 2006; 38:27-37; PMID:16311598; <http://dx.doi.org/10.1038/ng1679>
110. Luis TC, Naber BA, Roozen PP, Brugman MH, de Haas EF, Ghazvini M, et al. Canonical wnt signaling regulates hematopoiesis in a dosage-dependent fashion. *Cell Stem Cell* 2011; 9:345-56; PMID:21982234; <http://dx.doi.org/10.1016/j.stem.2011.07.017>
111. Pfützner E, Jähne R, Wissler M, Stoecklin E, Groner B. p300/CREB-binding protein enhances the prolactin-mediated transcriptional induction through direct interaction with the transactivation domain of Stat5, but does not participate in the Stat5-mediated suppression of the glucocorticoid response. *Mol Endocrinol* 1998; 12:1582-93; PMID:9773981; <http://dx.doi.org/10.1210/me.12.10.1582>
112. Gewinner C, Hart G, Zachara N, Cole R, Beisenherz-Huss C, Groner B. The coactivator of transcription CREB-binding protein interacts preferentially with the glycosylated form of Stat5. *J Biol Chem* 2004; 279:3563-72; PMID:14597631; <http://dx.doi.org/10.1074/jbc.M306449200>
113. Bakker WJ, van Dijk TB, Parren-van Amelsvoort M, Kolbus A, Yamamoto K, Steinlein P, et al. Differential regulation of Foxo3a target genes in erythropoiesis. *Mol Cell Biol* 2007; 27:3839-54; PMID:17353275; <http://dx.doi.org/10.1128/MCB.01662-06>
114. Kung AL, Rebel VI, Bronson RT, Ch'ng LE, Sieff CA, Livingston DM, et al. Gene dose-dependent control of hematopoiesis and hematologic tumor suppression by CBP. *Genes Dev* 2000; 14:272-7; PMID:10673499
115. Rebel VI, Kung AL, Tanner EA, Yang H, Bronson RT, Livingston DM. Distinct roles for CREB-binding protein and p300 in hematopoietic stem cell self-renewal. *Proc Natl Acad Sci U S A* 2002; 99:14789-94; PMID:12397173; <http://dx.doi.org/10.1073/pnas.232568499>
116. Chan WJ, Hannah RL, Dawson MA, Pridans C, Foster D, Joshi A, et al. The transcriptional coactivator Cbp regulates self-renewal and differentiation in adult hematopoietic stem cells. *Mol Cell Biol* 2011; 31:5046-60; PMID:22006020; <http://dx.doi.org/10.1128/MCB.05830-11>
117. Kranc KR, Schepers H, Rodrigues NP, Bamforth S, Villadsen E, Ferry H, et al. Cited2 is an essential regulator of adult hematopoietic stem cells. *Cell Stem Cell* 2009; 5:659-65; PMID:19951693; <http://dx.doi.org/10.1016/j.stem.2009.11.001>
118. Ikezoe T, Yang J, Nishioka C, Kojima S, Takeuchi A, Phillip Koeffler H, et al. Inhibition of signal transducer and activator of transcription 5 by the inhibitor of janus kinases stimulates dormant human leukemia CD34+ /CD38- cells and sensitizes them to anti-leukemia agents. *Int J Cancer* 2011; 128:2317-25; PMID:21128225; <http://dx.doi.org/10.1002/ijc.25806>
119. Cheng T, Rodrigues N, Shen H, Yang Y, Dombkowski D, Sykes M, et al. Hematopoietic stem cell quiescence maintained by p21cip1/waf1. *Science* 2000; 287:1804-8; PMID:10710306; <http://dx.doi.org/10.1126/science.287.5459.1804>
120. Yu JH, Zhu BM, Wickre M, Riedlinger G, Chen W, Hosui A, et al. The transcription factors signal transducer and activator of transcription 5A (STAT5A) and STAT5B negatively regulate cell proliferation through the activation of cyclin-dependent kinase inhibitor 2b (Cdkn2b) and Cdkn1a expression. *Hepatology* 2010; 52:1808-18; PMID:21038417; <http://dx.doi.org/10.1002/hep.23882>
121. Viale A, De Franco F, Orleth A, Cambiaghi V, Giuliani V, Bossi D, et al. Cell-cycle restriction limits DNA damage and maintains self-renewal of leukaemia stem cells. *Nature* 2009; 457:51-6; PMID:19122635; <http://dx.doi.org/10.1038/nature07618>

122. Kaluz S, Kaluzová M, Stanbridge EJ. Regulation of gene expression by hypoxia: integration of the HIF-transduced hypoxic signal at the hypoxia-responsive element. *Clin Chim Acta* 2008; 395:6-13; PMID:18505681; <http://dx.doi.org/10.1016/j.cca.2008.05.002>
123. Takubo K, Goda N, Yamada W, Iriuchishima H, Ikeda E, Kubota Y, et al. Regulation of the HIF-1 α level is essential for hematopoietic stem cells. *Cell Stem Cell* 2010; 7:391-402; PMID:20804974; <http://dx.doi.org/10.1016/j.stem.2010.06.020>
124. Simsek T, Kocabas F, Zheng J, Deberardinis RJ, Mahmoud AI, Olson EN, et al. The distinct metabolic profile of hematopoietic stem cells reflects their location in a hypoxic niche. *Cell Stem Cell* 2010; 7:380-90; PMID:20804973; <http://dx.doi.org/10.1016/j.stem.2010.07.011>
125. Calvi LM, Adams GB, Weibrecht KW, Weber JM, Olson DP, Knight MC, et al. Osteoblastic cells regulate the haematopoietic stem cell niche. *Nature* 2003; 425:841-6; PMID:14574413; <http://dx.doi.org/10.1038/nature02040>
126. Rizo A, Vellenga E, de Haan G, Schuringa JJ. Signaling pathways in self-renewing hematopoietic and leukemic stem cells: do all stem cells need a niche? *Hum Mol Genet* 2006; 15(Spec No 2):R210-9; PMID:16987886; <http://dx.doi.org/10.1093/hmg/ddl175>
127. Schofield R. The relationship between the spleen colony-forming cell and the haemopoietic stem cell. *Blood Cells* 1978; 4:7-25; PMID:747780
128. Zhang J, Niu C, Ye L, Huang H, He X, Tong WG, et al. Identification of the haematopoietic stem cell niche and control of the niche size. *Nature* 2003; 425:836-41; PMID:14574412; <http://dx.doi.org/10.1038/nature02041>
129. Rizo A, Dontje B, Vellenga E, de Haan G, Schuringa JJ. Long-term maintenance of human hematopoietic stem/progenitor cells by expression of BMI1. *Blood* 2008; 111:2621-30; PMID:18156489; <http://dx.doi.org/10.1182/blood-2007-08-106666>
130. Fatrai S, Schepers H, Tadema H, Vellenga E, Daenen SM, Schuringa JJ. Mucin1 expression is enriched in the human stem cell fraction of cord blood and is upregulated in majority of the AML cases. *Exp Hematol* 2008; 36:1254-65; PMID:18640764; <http://dx.doi.org/10.1016/j.exphem.2008.04.015>
131. Ahmad R, Raina D, Trivedi V, Ren J, Rajabi H, Kharbanda S, et al. MUC1 oncoprotein activates the I κ B kinase beta complex and constitutive NF- κ B signalling. *Nat Cell Biol* 2007; 9:1419-27; PMID:18037881; <http://dx.doi.org/10.1038/ncb1661>
132. Huang L, Ren J, Chen D, Li Y, Kharbanda S, Kufe D. MUC1 cytoplasmic domain coactivates Wnt target gene transcription and confers transformation. *Cancer Biol Ther* 2003; 2:700-4; PMID:14688481; <http://dx.doi.org/10.4161/cbt.2.6.610>
133. Huang L, Chen D, Liu D, Yin L, Kharbanda S, Kufe D. MUC1 oncoprotein blocks glycogen synthase kinase 3 β -mediated phosphorylation and degradation of beta-catenin. *Cancer Res* 2005; 65:10413-22; PMID:16288032; <http://dx.doi.org/10.1158/0008-5472.CAN-05-2474>
134. Wen Y, Caffrey TC, Wheelock MJ, Johnson KR, Hollingsworth MA. Nuclear association of the cytoplasmic tail of MUC1 and beta-catenin. *J Biol Chem* 2003; 278:38029-39; PMID:12832415; <http://dx.doi.org/10.1074/jbc.M304333200>
135. Wei X, Xu H, Kufe D. Human MUC1 oncoprotein regulates p53-responsive gene transcription in the genotoxic stress response. *Cancer Cell* 2005; 7:167-78; PMID:15710329; <http://dx.doi.org/10.1016/j.ccr.2005.01.008>
136. Rahn JJ, Chow JW, Horne GJ, Mah BK, Emerman JT, Hoffman P, et al. MUC1 mediates transendothelial migration in vitro by ligating endothelial cell ICAM-1. *Clin Exp Metastasis* 2005; 22:475-83; PMID:16320110; <http://dx.doi.org/10.1007/s10585-005-3098-x>
137. Rahn JJ, Shen Q, Mah BK, Hugh JC. MUC1 initiates a calcium signal after ligation by intercellular adhesion molecule-1. *J Biol Chem* 2004; 279:29386-90; PMID:15169768; <http://dx.doi.org/10.1074/jbc.C400010200>
138. Adams GB, Chabner KT, Alley IR, Olson DP, Szczepiorkowski ZM, Poznansky MC, et al. Stem cell engraftment at the endosteal niche is specified by the calcium-sensing receptor. *Nature* 2006; 439:599-603; PMID:16382241; <http://dx.doi.org/10.1038/nature04247>

© 2012 Landes Bioscience.

Do not distribute.